

NDA 20-287/S-018

Pharmacia & Upjohn Company  
Attention: Ms. Leslie A. Franks  
7000 Portage Road  
Kalamazoo, Michigan 49001

AUG 3 2000

Dear Ms. Franks:

Please refer to your supplemental new drug application dated February 20, 2000, received March 1, 2000, submitted under section 505(b) of the Federal Food, Drug, and Cosmetic Act for Fragmin® (dalteparin sodium injection).

We acknowledge receipt of your submission dated July 21, 2000.

This supplemental new drug application provides for the addition of a postoperative dosing regimen for Fragmin; and (2) content changes to the package insert including the CLINICAL TRIALS section, "Hip Replacement Surgery" subsection; the ADVERSE REACTIONS section, "Hip Replacement Surgery" subsection; and the DOSAGE AND ADMINISTRATION section, "Hip Replacement Surgery" subsection.

We have completed the review of this supplemental application, as amended, and have concluded that adequate information has been presented to demonstrate that the drug product is safe and effective for use as recommended in the agreed upon labeling text. Accordingly, the supplemental application is approved effective on the date of this letter.

The final printed labeling (FPL) must be identical to the enclosed labeling text (text for package insert).

Please submit 20 paper copies of the FPL as soon as it is available, in no case more than 30 days after it is printed. Please individually mount ten of the copies on heavy-weight paper or similar material. Alternatively, you may submit the FPL electronically according to the guidance for industry titled *Providing Regulatory Submissions in Electronic Format NDAs* (January 1999). For administrative purposes, this submission should be designated "FPL for approved supplement NDA 20-287/S-018." Approval of this submission by FDA is not required before the labeling is used.

Be advised that, as of April 1, 1999, all applications for new active ingredients, new dosage forms, new indications, new routes of administration, and new dosing regimens are required to contain an assessment of the safety and effectiveness of the product in pediatric patients unless this requirement

is waived or deferred (63 *FR* 66632). We note that you have not fulfilled the requirements of 21 CFR 314.55 (or 601.27). We are deferring submission of your pediatric studies until December 2, 2000. However, in the interim, please submit your pediatric drug development plans within 120 days from the date of this letter unless you believe a waiver is appropriate. Within approximately 120 days of receipt of your pediatric drug development plan, we will review your plan and notify you of its adequacy. If you believe that this drug qualifies for a waiver of the pediatric study requirement, you should submit a request for a waiver with supporting information and documentation in accordance with the provisions of 21 CFR 314.55 within 60 days from the date of this letter. We will notify you within 120 days of receipt of your response whether a waiver is granted. If a waiver is not granted, we will ask you to submit your pediatric drug development plans within 120 days from the date of denial of the waiver.

Pediatric studies conducted under the terms of section S05A of the Federal Food, Drug, and Cosmetic Act may result in additional marketing exclusivity for certain products (pediatric exclusivity). You should refer to the *Guidance for Industry on Qualifying for Pediatric Exclusivity* (available on our web site at [www.fda.gov/cder/pediatric](http://www.fda.gov/cder/pediatric)) for details. If you wish to qualify for pediatric exclusivity you should submit a "Proposed Pediatric Study Request" (PPSR) in addition to your plans for pediatric drug development described above. We recommend that you submit a Proposed Pediatric Study Request within 120 days from the date of this letter. If you are unable to meet this time frame but are interested in pediatric exclusivity, please notify the division in writing. FDA generally will not accept studies submitted to an NDA before issuance of a Written Request as responsive to a Written Request. Sponsors should obtain a Written Request before submitting pediatric studies to an NDA. If you do not submit a PPSR or indicate that you are interested in pediatric exclusivity, we will review your pediatric drug development plan and notify you of its adequacy. Please note that satisfaction of the requirements in 21 CFR 314.55 alone may not qualify you for pediatric exclusivity. FDA does not necessarily ask a sponsor to complete the same scope of studies to qualify for pediatric exclusivity as it does to fulfill the requirements of the pediatric rule.

In addition, please submit three copies of the introductory promotional materials that you propose to use for this product. All proposed materials should be submitted in draft or mock-up form, not final print. Please submit one copy to this Division and two copies of both the promotional materials and the package insert directly to:

Division of Drug Marketing, Advertising, and Communications, HFD-42  
Food and Drug Administration  
5600 Fishers Lane  
Rockville, Maryland 20857

If a letter communicating important information about this drug product (i.e., a “Dear Health Care Practitioner” letter) is issued to physicians and others responsible for patient care, we request that you submit a copy of the letter to this NDA and a copy to the following address:

MEDWATCH, HF-2  
FDA  
5600 Fishers Lane  
Rockville,MD 20857

Please submit one market package of the drug product when it is available.

We remind you that you must comply with the requirements for an approved NDA set forth under 21 CFR 314.80 and 314.81.

If you have any questions, call Karen Oliver, Regulatory Project Manager, at (301) 827-7457.

Sincerely,

Lilia Talarico, M.D.  
Director  
Division of Gastrointestinal and Coagulation Drug  
Products  
Office of Drug Evaluation III  
Center for Drug Evaluation and Research



## dalteparin sodium injection



Pharmacia  
& Upjohn

For *Subcutaneous* Use Only

### SPINAL/EPIDURAL HEMATOMAS

When neuraxial anesthesia (epidural/spinal anesthesia) or spinal puncture is employed, patients anticoagulated or scheduled to be anticoagulated with low molecular weight heparins or heparinoids for prevention of thromboembolic complications are at risk of developing an epidural or spinal hematoma which can result in long-term or permanent paralysis.

The risk of these events is increased by the use of indwelling epidural catheters for administration of analgesia or by the concomitant use of drugs affecting hemostasis such as non steroidal anti-inflammatory drugs (NSAIDs), platelet inhibitors, or other anticoagulants. The risk also appears to be increased by traumatic or repeated epidural or spinal puncture.

Patients should be frequently monitored for signs and symptoms of neurological impairment. If neurological compromise is noted, urgent treatment is necessary.

The physician should consider the potential benefit versus risk before neuraxial intervention in patients anticoagulated or to be anticoagulated for thromboprophylaxis (also see WARNINGS, Hemorrhage and PRECAUTIONS, Drug Interactions).

### DESCRIPTION


FRAGMIN Injection (dalteparin sodium injection) is a sterile, low molecular weight heparin. It is available in single-dose, prefilled syringes and a multiple-dose vial. With reference to the W.H.O. First International Low Molecular Weight Heparin Reference Standard, each syringe contains 2500 (16 mg dalteparin sodium) or 5000 (32 mg dalteparin sodium) anti-Factor Xa international units (IU) in 0.2 mL. Each 9.5 mL vial contains 10,000 (64 mg dalteparin sodium) anti-Factor Xa IU per 1 mL, for a total of 95,000 anti-Factor Xa IU per vial.

Each prefilled syringe also contains Water for Injection and sodium chloride, when required, to maintain physiologic ionic strength. The prefilled syringes are preservative free. Each multiple-dose vial also contains Water for Injection and 14 mg of benzyl alcohol per mL as a preservative. The pH of both formulations is 5.0 to 7.5.

Dalteparin sodium is produced through controlled nitrous acid depolymerization of sodium heparin from porcine intestinal mucosa followed by a chromatographic purification process. It is composed of strongly acidic sulphated polysaccharide chains (oligosaccharide, containing 2,5-anhydro-D-mannitol residues as end groups) with an average molecular weight of 5000 and about 90% of the material within the range 2000–9000. The molecular weight distribution is:

< 3000 daltons	3.0–15.0%
3000 to 8000 daltons	65.0–78.0%
> 8000 daltons	14.0–26.0%

### Structural Formula



R = H or SO<sub>3</sub>Na  
R<sub>1</sub> = COCH<sub>3</sub> or SO<sub>3</sub>Na  
R<sub>2</sub> = H    R<sub>3</sub> = COONa  
          or  
R<sub>2</sub> = COONa    R<sub>3</sub> = H

n = 3–20

# Fragmin

brand of dalteparin sodium injection

## CLINICAL PHARMACOLOGY

Dalteparin is a low molecular weight heparin with antithrombotic properties. It acts by enhancing the inhibition of Factor Xa and thrombin by antithrombin. In man, dalteparin potentiates preferentially the inhibition of coagulation Factor Xa, while only slightly affecting clotting time, e.g., activated partial thromboplastin time (APTT).

### Pharmacodynamics:

Doses of FRAGMIN Injection of up to 10,000 anti-Factor Xa IU administered subcutaneously as a single dose or two 5000 IU doses 12 hours apart to healthy subjects do not produce a significant change in platelet aggregation, fibrinolysis, or global clotting tests such as prothrombin time (PT), thrombin time (TT) or APTT. Subcutaneous (s.c.) administration of doses of 5000 IU bid of FRAGMIN for seven consecutive days to patients undergoing abdominal surgery did not markedly affect APTT, Platelet Factor 4 (PF4), or lipoprotein lipase.

### Pharmacokinetics:

Mean peak levels of plasma anti-Factor Xa activity following single s.c. doses of 2500, 5000 and 10,000 IU were  $0.19 \pm 0.04$ ,  $0.41 \pm 0.07$  and  $0.82 \pm 0.10$  IU/mL, respectively, and were attained in about 4 hours in most subjects. Absolute bioavailability in healthy volunteers, measured as the anti-Factor Xa activity, was  $87 \pm 6\%$ . Increasing the dose from 2500 to 10,000 IU resulted in an overall increase in anti-Factor Xa AUC that was greater than proportional by about one-third.

Peak anti-Factor Xa activity increased more or less linearly with dose over the same dose range. There appeared to be no appreciable accumulation of anti-Factor Xa activity with twice-daily dosing of 100 IU/kg s.c. for up to 7 days.

The volume of distribution for dalteparin anti-Factor Xa activity was 40 to 60 mL/kg. The mean plasma clearances of dalteparin anti-Factor Xa activity in normal volunteers following single intravenous bolus doses of 30 and 120 anti-Factor Xa IU/kg were  $24.6 \pm 5.4$  and  $15.6 \pm 2.4$  mL/hr/kg, respectively. The corresponding mean disposition half-lives are  $1.47 \pm 0.3$  and  $2.5 \pm 0.3$  hours.

Following intravenous doses of 40 and 60 IU/kg, mean terminal half-lives were  $2.1 \pm 0.3$  and  $2.3 \pm 0.4$  hours, respectively. Longer apparent terminal half-lives (3 to 5 hours) are observed following s.c. dosing, possibly due to delayed absorption. In patients with chronic renal insufficiency requiring hemodialysis, the mean terminal half-life of anti-Factor Xa activity following a single intravenous dose of 5000 IU FRAGMIN was  $5.7 \pm 2.0$  hours, i.e. considerably longer than values observed in healthy volunteers, therefore, greater accumulation can be expected in these patients.

## CLINICAL TRIALS

### Prophylaxis of Ischemic Complications in Unstable Angina and Non-Q-Wave Myocardial Infarction:

In a double-blind, randomized, placebo-controlled clinical trial, patients who recently experienced unstable angina with EKG changes or non-Q-wave myocardial infarction (MI) were randomized to FRAGMIN Injection 120 IU/kg every 12 hours subcutaneously (s.c.) or placebo every 12 hours s.c. In this trial, unstable angina was defined to include only angina with EKG changes. All patients, except when contraindicated, were treated concurrently with aspirin (75 mg once daily) and beta blockers. Treatment was initiated within 72 hours of the event (the majority of patients received treatment within 24 hours) and continued for 5 to 8 days. A total of 1506 patients were enrolled and treated; 746 received FRAGMIN and 760 received placebo. The mean age of the study population was 68 years (range 40 to 90 years) and the majority of patients were white (99.7%) and male (63.9%). The combined incidence of the double endpoint of death or myocardial infarction was lower for FRAGMIN compared with placebo at 6 days after initiation of therapy. These results were observed in an analysis of all-randomized and all-treated patients. The combined incidence of death, MI, need for intravenous (i.v.) heparin or i.v. nitroglycerin, and revascularization was also lower for FRAGMIN than for placebo (see Table 1).

# Fragmin

brand of dalteparin sodium injection

**Table 1**  
**Efficacy of FRAGMIN in the Prophylaxis of Ischemic Complications in Unstable Angina and Non-Q-Wave Myocardial Infarction**

Indication	Dosing Regimen	
	<b>FRAGMIN</b> 120 IU/kg/12 hr s.c.	<b>Placebo</b> q 12 hr s.c.
All Treated Unstable Angina and Non-Q-Wave MI Patients	746	760
Primary Endpoints – 6 day timepoint Death, MI	13/741 (1.8%) <sup>1</sup>	36/757 (4.8%)
Secondary Endpoints – 6 day timepoint Death, MI, i.v. heparin, i.v. nitroglycerin, Revascularization	59/739 (8.0%) <sup>1</sup>	106/756 (14.0%)

<sup>1</sup> p-value = 0.001

In a second randomized, controlled trial designed to evaluate long-term treatment with FRAGMIN (days 6 to 45), data were also collected comparing 1-week (5 to 8 days) treatment of FRAGMIN 120 IU/kg every 12 hours s.c. with heparin at an APTT-adjusted dosage. All patients, except when contraindicated, were treated concurrently with aspirin (100 to 165 mg per day). Of the total enrolled study population of 1499 patients, 1482 patients were treated; 751 received FRAGMIN and 731 received heparin. The mean age of the study population was 64 years (range 25 to 92 years) and the majority of patients were white (96.0%) and male (64.2%). The incidence of the combined triple endpoint of death, myocardial infarction, or recurrent angina during this 1-week treatment period (5 to 8 days) was 9.3% for FRAGMIN and 7.6% for heparin (p=0.323).

## **Prophylaxis of Deep Vein Thrombosis in Patients Following Hip Replacement Surgery:**

In an open-label randomized study, FRAGMIN 5000 IU administered once daily s.c. was compared with warfarin sodium, administered orally, in patients undergoing hip replacement surgery. Treatment with FRAGMIN was initiated with a 2500 IU dose s.c. within 2 hours before surgery, followed by a 2500 IU dose s.c. the evening of the day of surgery. Then, a dosing regimen of FRAGMIN 5000 IU s.c. once daily was initiated on the first postoperative day. The first dose of warfarin sodium was given the evening before surgery, then continued daily at a dose adjusted for INR 2.0 to 3.0. Treatment in both groups was then continued for 5 to 9 days postoperatively. Of the total enrolled study population of 580 patients, 553 were treated and 550 underwent surgery. Of those who underwent surgery, 271 received FRAGMIN and 279 received warfarin sodium. The mean age of the study population was 63 years (range 20 to 92 years) and the majority of patients were white (91.1%) and female (52.9%). The incidence of deep vein thrombosis (DVT), any vein, as determined by evaluable venography, was significantly lower for the group treated with FRAGMIN compared with patients treated with warfarin sodium (28/192 vs 49/190; p=0.006) (see Table 2).

**Table 2**  
**Efficacy of FRAGMIN in the Prophylaxis of Deep Vein Thrombosis Following Hip Replacement Surgery**

Indication	Dosing Regimen	
	<b>FRAGMIN</b> 5000 IU qd <sup>1</sup> s.c.	<b>Warfarin Sodium</b> qd <sup>2</sup> oral
All Treated Hip Replacement Surgery Patients	271	279
Treatment Failures in Evaluable Patients		
DVT, Total	28/192 (14.6%) <sup>3</sup>	49/190 (25.8%)
Proximal DVT	10/192 (5.2%) <sup>4</sup>	16/190 (8.4%)
PE	2/271 (0.7%)	2/279 (0.7%)

<sup>1</sup> The daily dose on the day of surgery was divided: 2500 IU was given two hours before surgery and again in the evening of the day of surgery.

<sup>2</sup> Warfarin sodium dosage was adjusted to maintain a prothrombin time index of 1.4 to 1.5, corresponding to an International Normalized Ratio (INR) of approximately 2.5.

<sup>3</sup> p-value = 0.006

<sup>4</sup> p-value = 0.185

# Fragmin

brand of dalteparin sodium injection

In a second single-center, double-blind study of patients undergoing hip replacement surgery, FRAGMIN 5000 IU once daily s.c. starting the evening before surgery, was compared with heparin 5000 IU s.c. tid, starting the morning of surgery. Treatment in both groups was continued for up to 9 days postoperatively. Of the total enrolled study population of 140 patients, 139 were treated and 136 underwent surgery. Of those who underwent surgery, 67 received FRAGMIN and 69 received heparin. The mean age of the study population was 69 years (range 42 to 87 years) and the majority of patients were female (58.8%). In the intent-to-treat analysis, the incidence of proximal DVT was significantly lower for patients treated with FRAGMIN compared with patients treated with heparin (6/67 vs 18/69; p=0.012). Further, the incidence of pulmonary embolism detected by lung scan was also significantly lower in the group treated with FRAGMIN (9/67 vs 19/69; p=0.032).

A third multi-center, double-blind, randomized study evaluated a postoperative dosing regimen of FRAGMIN for thromboprophylaxis following total hip replacement surgery. Patients received either FRAGMIN or warfarin sodium, randomized into one of three treatment groups. One group of patients received the first dose of FRAGMIN 2500 IU s.c. within 2 hours before surgery, followed by another dose of FRAGMIN 2500 IU s.c. at least 4 hours (6.6 ± 2.3 hr) after surgery. Another group received the first dose of FRAGMIN 2500 IU s.c. at least 4 hours (6.6 ± 2.4 hr) after surgery. Then, both of these groups began a dosing regimen of FRAGMIN 5000 IU once daily s.c. on postoperative day 1. The third group of patients received warfarin sodium the evening of the day of surgery, then continued daily at a dose adjusted for INR 2.0 to 3.0. Treatment for all groups was continued for 4 to 8 days postoperatively, after which time all patients underwent bilateral venography.

In the total enrolled study population of 1501 patients, 1472 patients were treated; 496 received FRAGMIN (first dose before surgery), 487 received FRAGMIN (first dose after surgery) and 489 received warfarin sodium. The mean age of the study population was 63 years (range 18 to 91 years) and the majority of patients were white (94.4%) and female (51.8%).

Administration of the first dose of FRAGMIN after surgery was as effective in reducing the incidence of thromboembolic events as administration of the first dose of FRAGMIN before surgery (44/336 vs 37/338; p=0.448). Both dosing regimens of FRAGMIN were more effective than warfarin sodium in reducing the incidence of thromboembolic events following hip replacement surgery.

**Prophylaxis of Deep Vein Thrombosis Following Abdominal Surgery in Patients at Risk for Thromboembolic Complications:**

Abdominal surgery patients at risk include those who are over 40 years of age, obese, undergoing surgery under general anesthesia lasting longer than 30 minutes, or who have additional risk factors such as malignancy or a history of deep vein thrombosis or pulmonary embolism.

FRAGMIN administered once daily s.c. beginning prior to surgery and continuing for 5 to 10 days after surgery, was shown to reduce the risk of DVT in patients at risk for thromboembolic complications in two double-blind, randomized, controlled clinical trials performed in patients undergoing major abdominal surgery. In the first study, a total of 204 patients were enrolled and treated; 102 received FRAGMIN and 102 received placebo. The mean age of the study population was 64 years (range 40 to 98 years) and the majority of patients were female (54.9%). In the second study, a total of 391 patients were enrolled and treated; 195 received FRAGMIN and 196 received heparin. The mean age of the study population was 59 years (range 30 to 88 years) and the majority of patients were female (51.9%). As summarized in the following tables, FRAGMIN 2500 IU was superior to placebo and similar to heparin in reducing the risk of DVT (see Tables 3 and 4).

**Table 3**  
**Efficacy of FRAGMIN in the Prophylaxis of Deep Vein Thrombosis Following Abdominal Surgery**

Indication	Dosing Regimen	
	FRAGMIN 2500 IU qd s.c.	Placebo qd s.c.
All Treated Abdominal Surgery Patients	102	102
Treatment Failures in Evaluable Patients		
Total Thromboembolic Events	4/91 (4.4%) <sup>1</sup>	16/91 (17.6%)
Proximal DVT	0	5/91 (5.5%)
Distal DVT	4/91 (4.4%)	11/91 (12.1%)
PE	0	2/91 (2.2%) <sup>2</sup>

<sup>1</sup> p-value = 0.008

<sup>2</sup> Both patients also had DVT, 1 proximal and 1 distal

# Fragmin

brand of dalteparin sodium injection

**Table 4**  
**Efficacy of FRAGMIN in the Prophylaxis of Deep Vein Thrombosis Following Abdominal Surgery**

Indication	Dosing Regimen	
	FRAGMIN 2500 IU qd s.c.	Heparin 5000 U bid s.c.
All Treated Abdominal Surgery Patients	195	196
Treatment Failures in Evaluable Patients		
Total Thromboembolic Events	7/178 (3.9%) <sup>1</sup>	7/174 (4.0%)
Proximal DVT	3/178 (1.7%)	4/174 (2.3%)
Distal DVT	3/178 (1.7%)	3/174 (1.7%)
PE	1/178 (0.6%)	0

<sup>1</sup> p-value = 0.74

In a third double-blind, randomized study performed in patients undergoing major abdominal surgery with malignancy, FRAGMIN 5000 IU once daily was compared with FRAGMIN 2500 IU once daily. Treatment was continued for 6 to 8 days. A total of 1375 patients were enrolled and treated; 679 received FRAGMIN 5000 IU and 696 received 2500 IU. The mean age of the combined groups was 71 years (range 40 to 95 years). The majority of patients were female (51.0%). The study showed that FRAGMIN 5000 IU once daily was more effective than FRAGMIN 2500 IU once daily in reducing the risk of DVT in patients undergoing abdominal surgery with malignancy (see Table 5).

**Table 5**  
**Efficacy of FRAGMIN in the Prophylaxis of Deep Vein Thrombosis Following Abdominal Surgery**

Indication	Dosing Regimen	
	FRAGMIN 2500 IU qd s.c.	FRAGMIN 5000 IU qd s.c.
All Treated Abdominal Surgery Patients <sup>1</sup>	696	679
Treatment Failures in Evaluable Patients		
Total Thromboembolic Events	99/656 (15.1%) <sup>2</sup>	60/645 (9.3%)
Proximal DVT	18/657 (2.7%)	14/646 (2.2%)
Distal DVT	80/657 (12.2%)	41/646 (6.3%)
PE		
Fatal	1/674 (0.1%)	1/669 (0.1%)
Non-fatal	2	4

<sup>1</sup> Major abdominal surgery with malignancy

<sup>2</sup> p-value = 0.001

## INDICATIONS AND USAGE

FRAGMIN injection is indicated for the prophylaxis of ischemic complications in unstable angina and non-Q-wave myocardial infarction, when concurrently administered with aspirin therapy (as described in CLINICAL TRIALS, Prophylaxis of Ischemic Complications in Unstable Angina and Non-Q-Wave Myocardial Infarction).

FRAGMIN is also indicated for the prophylaxis of deep vein thrombosis (DVT), which may lead to pulmonary embolism (PE):

- In patients undergoing hip replacement surgery;
- In patients undergoing abdominal surgery who are at risk for thromboembolic complications.

## CONTRAINDICATIONS

FRAGMIN Injection is contraindicated in patients with known hypersensitivity to the drug, active major bleeding, or thrombocytopenia associated with positive *in vitro* tests for anti-platelet antibody in the presence of FRAGMIN.

Patients undergoing regional anesthesia should not receive FRAGMIN for unstable angina or non-Q-wave myocardial infarction due to an increased risk of bleeding associated with the dosage of FRAGMIN recommended for unstable angina and non-Q-wave myocardial infarction.



# Fragmin

## brand of dalteparin sodium injection

Patients with known hypersensitivity to heparin or pork products should not be treated with FRAGMIN.

### **WARNINGS**

FRAGMIN Injection is not intended for intramuscular administration.

FRAGMIN cannot be used interchangeably (unit for unit) with unfractionated heparin or other low molecular weight heparins.

**FRAGMIN should be used with extreme caution in patients with history of heparin-induced thrombocytopenia.**

#### **Hemorrhage:**

FRAGMIN, like other anticoagulants, should be used with extreme caution in patients who have an increased risk of hemorrhage, such as those with severe uncontrolled hypertension, bacterial endocarditis, congenital or acquired bleeding disorders, active ulceration and angiodysplastic gastrointestinal disease, hemorrhagic stroke, or shortly after brain, spinal or ophthalmological surgery.

**Spinal or epidural hematomas can occur with the associated use of low molecular weight heparins or heparinoids and neuraxial (spinal/epidural) anesthesia or spinal puncture, which can result in long-term or permanent paralysis. The risk of these events is higher with the use of indwelling epidural catheters or concomitant use of additional drugs affecting hemostasis such as NSAIDs (see boxed WARNING and ADVERSE REACTIONS, Ongoing Safety Surveillance).**

As with other anticoagulants, bleeding can occur at any site during therapy with FRAGMIN. An unexpected drop in hematocrit or blood pressure should lead to a search for a bleeding site.

#### **Thrombocytopenia:**

In clinical trials, thrombocytopenia with platelet counts of  $<100,000/\text{mm}^3$  and  $<50,000/\text{mm}^3$  occurred in  $<1\%$  and  $<1\%$ , respectively. In clinical practice, rare cases of thrombocytopenia with thrombosis have also been observed.

Thrombocytopenia of any degree should be monitored closely. Heparin-induced thrombocytopenia can occur with the administration of FRAGMIN. The incidence of this complication is unknown at present.

#### **Miscellaneous:**

The multiple-dose vial of FRAGMIN contains benzyl alcohol as a preservative. Benzyl alcohol has been reported to be associated with a fatal "Gasping Syndrome" in premature infants. Because benzyl alcohol may cross the placenta, FRAGMIN preserved with benzyl alcohol should not be used in pregnant women (see PRECAUTIONS, Pregnancy Category B., Nonteratogenic Effects).

### **PRECAUTIONS**

#### **General:**

FRAGMIN Injection should not be mixed with other injections or infusions unless specific compatibility data are available that support such mixing.

FRAGMIN should be used with caution in patients with bleeding diathesis, thrombocytopenia or platelet defects; severe liver or kidney insufficiency, hypertensive or diabetic retinopathy, and recent gastrointestinal bleeding.

If a thromboembolic event should occur despite dalteparin prophylaxis, FRAGMIN should be discontinued and appropriate therapy initiated.

#### **Drug Interactions:**

FRAGMIN should be used with care in patients receiving oral anticoagulants, platelet inhibitors, and thrombolytic agents because of increased risk of bleeding (see PRECAUTIONS, Laboratory Tests). Aspirin, unless contraindicated, is recommended in patients treated for unstable angina or non-Q-wave myocardial infarction (see DOSAGE AND ADMINISTRATION).

#### **Laboratory Tests:**

Periodic routine complete blood counts, including platelet count, and stool occult blood tests are recommended during the course of treatment with FRAGMIN. No special monitoring of blood clotting times (e.g., APTT) is needed.

When administered at recommended prophylaxis doses, routine coagulation tests such as Prothrombin Time (PT) and Activated Partial Thromboplastin Time (APTT) are relatively insensitive measures of FRAGMIN activity and, therefore, unsuitable for monitoring.

# Fragmin

brand of dalteparin sodium injection

## Drug/Laboratory Test Interactions:

### *Elevations of Serum Transaminases:*

Asymptomatic increases in transaminase levels (SGOT/AST and SGPT/ALT) greater than three times the upper limit of normal of the laboratory reference range have been reported in 1.7% and 4.3%, respectively, of patients during treatment with FRAGMIN. Similar significant increases in transaminase levels have also been observed in patients treated with heparin and other low molecular weight heparins. Such elevations are fully reversible and are rarely associated with increases in bilirubin. Since transaminase determinations are important in the differential diagnosis of myocardial infarction, liver disease and pulmonary emboli, elevations that might be caused by drugs like FRAGMIN should be interpreted with caution.

## Carcinogenicity, Mutagenesis, Impairment of Fertility:

Dalteparin sodium has not been tested for its carcinogenic potential in long-term animal studies. It was not mutagenic in the *in vitro* Ames Test, mouse lymphoma cell forward mutation test and human lymphocyte chromosomal aberration test and in the *in vivo* mouse micronucleus test. Dalteparin sodium at subcutaneous doses up to 1200 IU/kg (7080 IU/m<sup>2</sup>) did not affect the fertility or reproductive performance of male and female rats.

## Pregnancy: Pregnancy Category B.

### Teratogenic Effects:

Reproduction studies with dalteparin sodium at intravenous doses up to 2400 IU/kg (14,160 IU/m<sup>2</sup>) in pregnant rats and 4800 IU/kg (40,800 IU/m<sup>2</sup>) in pregnant rabbits did not produce any evidence of impaired fertility or harm to the fetuses. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

### Nonteratogenic Effects:

Cases of "Gasping Syndrome" have occurred when large amounts of benzyl alcohol have been administered (99–404 mg/kg/day). The 9.5 mL multiple-dose vial of FRAGMIN contains 14 mg/mL of benzyl alcohol.

### Nursing Mothers:

It is not known whether dalteparin sodium is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when FRAGMIN is administered to a nursing mother.

### Pediatric Use:

Safety and effectiveness in pediatric patients have not been established.

## ADVERSE REACTIONS

### Hemorrhage:

The incidence of hemorrhagic complications during treatment with FRAGMIN Injection has been low. The most commonly reported side effect is hematoma at the injection site. The incidence of bleeding may increase with higher doses; however, in abdominal surgery patients with malignancy, no significant increase in bleeding was observed when comparing FRAGMIN 5000 IU to either FRAGMIN 2500 IU or low dose heparin.

In a trial comparing FRAGMIN 5000 IU once daily to FRAGMIN 2500 IU once daily in patients undergoing surgery for malignancy, the incidence of bleeding events was 4.6% and 3.6%, respectively (n.s.). In a trial comparing FRAGMIN 5000 IU once daily to heparin 5000 U twice daily, the incidence of bleeding events was 3.2% and 2.7%, respectively (n.s.) in the malignancy subgroup.

### *Unstable Angina and Non-Q-Wave Myocardial Infarction:*

Table 6 summarizes major bleeding events that occurred with FRAGMIN, heparin, and placebo in clinical trials of unstable angina and non-Q-wave myocardial infarction.

# Fragmin

brand of dalteparin sodium injection

**Table 6**  
**Major Bleeding Events in Unstable Angina and Non-Q-Wave Myocardial Infarction**

Indication	Dosing Regimen		
Unstable Angina and Non-Q-Wave MI	<b>FRAGMIN</b> 120 IU/kg/12 hr s.c. <sup>1</sup>	<b>Heparin</b> i.v. and s.c. <sup>2</sup>	<b>Placebo</b> q 12 hr s.c.
Major Bleeding Events <sup>3,4</sup>	15/1497 (1.0%)	7/731 (1.0%)	4/760 (0.5%)

<sup>1</sup> Treatment was administered for 5 to 8 days.

<sup>2</sup> Heparin i.v. infusion for at least 48 hours, APPT 1.5 to 2 times control, then 12,500 U s.c. every 12 hours for 5 to 8 days.

<sup>3</sup> Aspirin (75 to 165 mg per day) and beta blocker therapies were administered concurrently.

<sup>4</sup> Bleeding events were considered major if: 1) accompanied by a decrease in hemoglobin of  $\geq 2$  g/dL in connection with clinical symptoms; 2) a transfusion was required; 3) bleeding led to interruption of treatment or death; or 4) intracranial bleeding.

## Hip Replacement Surgery:

Table 7 summarizes: 1) all major bleeding events and, 2) other bleeding events possibly or probably related to treatment with FRAGMIN (preoperative dosing regimen), warfarin sodium, or heparin in two hip replacement surgery clinical trials.

**Table 7**  
**Bleeding Events Following Hip Replacement Surgery**

Indication	FRAGMIN vs Warfarin Sodium		FRAGMIN vs Heparin	
	Dosing Regimen		Dosing Regimen	
	<b>FRAGMIN</b> 5000 IU qd s.c. (n=274 <sup>2</sup> )	<b>Warfarin</b> Sodium <sup>1</sup> oral (n=279)	<b>FRAGMIN</b> 5000 IU qd s.c. (n=69 <sup>4</sup> )	<b>Heparin</b> 5000 U tid s.c. (n=69)
Hip Replacement Surgery				
Major Bleeding Events <sup>3</sup>	7/274 (2.6%)	1/279 (0.4%)	0	3/69 (4.3%)
Other Bleeding Events <sup>5</sup>				
Hematuria	8/274 (2.9%)	5/279 (1.8%)	0	0
Wound Hematoma	6/274 (2.2%)	0	0	0
Injection Site Hematoma	3/274 (1.1%)	NA	2/69 (2.9%)	7/69 (10.1%)

<sup>1</sup> Warfarin sodium dosage was adjusted to maintain a prothrombin time index of 1.4 to 1.5, corresponding to an International Normalized Ratio (INR) of approximately 2.5.

<sup>2</sup> Includes three treated patients who did not undergo a surgical procedure.

<sup>3</sup> A bleeding event was considered major if: 1) hemorrhage caused a significant clinical event, 2) it was associated with a hemoglobin decrease of  $\geq 2$  g/dL or transfusion of 2 or more units of blood products, 3) it resulted in reoperation due to bleeding, or 4) it involved retroperitoneal or intracranial hemorrhage.

<sup>4</sup> Includes two treated patients who did not undergo a surgical procedure.

<sup>5</sup> Occurred at a rate of at least 2% in the group treated with FRAGMIN 5000 IU once daily.

Six of the patients treated with FRAGMIN experienced seven major bleeding events. Two of the events were wound hematoma (one requiring reoperation), three were bleeding from the operative site, one was intraoperative bleeding due to vessel damage, and one was gastrointestinal bleeding. None of the patients experienced retroperitoneal or intracranial hemorrhage nor died of bleeding complications.

In the third hip replacement surgery clinical trial, the incidence of major bleeding events was similar in all three treatment groups: 3.6% (18/496) for patients who started FRAGMIN before surgery; 2.5% (12/487) for patients who started FRAGMIN after surgery; and 3.1% (15/489) for patients treated with warfarin sodium.

## Abdominal Surgery:

Table 8 summarizes bleeding events that occurred in clinical trials which studied FRAGMIN 2500 and 5000 IU administered once daily to abdominal surgery patients.

# Fragmin

brand of dalteparin sodium injection

**Table 8**  
**Bleeding Events Following Abdominal Surgery**

Indication	FRAGMIN vs Heparin				FRAGMIN vs Placebo		FRAGMIN vs FRAGMIN	
	Dosing Regimen				Dosing Regimen		Dosing Regimen	
	FRAGMIN 2500 IU qd s.c.	Heparin 5000 U bid s.c.	FRAGMIN 5000 IU qd s.c.	Heparin 5000 U bid s.c.	FRAGMIN 2500 IU qd s.c.	Placebo qd s.c.	FRAGMIN 2500 IU qd s.c.	FRAGMIN 5000 IU qd s.c.
Abdominal Surgery								
Postoperative Transfusions	26/459 (5.7%)	36/454 (7.9%)	81/508 (15.9%)	63/498 (12.7%)	14/182 (7.7%)	13/182 (7.1%)	89/1025 (8.7%)	125/1033 (12.1%)
Wound Hematoma	16/467 (3.4%)	18/467 (3.9%)	12/508 (2.4%)	6/498 (1.2%)	2/79 (2.5%)	2/77 (2.6%)	1/1030 (0.1%)	4/1039 (0.4%)
Reoperation Due to Bleeding	2/392 (0.5%)	3/392 (0.8%)	4/508 (0.8%)	2/498 (0.4%)	1/79 (1.3%)	1/78 (1.3%)	2/1030 (0.2%)	13/1038 (1.3%)
Injection Site Hematoma	1/466 (0.2%)	5/464 (1.1%)	36/506 (7.1%)	47/493 (9.5%)	8/172 (4.7%)	2/174 (1.1%)	36/1026 (3.5%)	57/1035 (5.5%)

**Thrombocytopenia: See WARNINGS: Thrombocytopenia.**

**Other:**

## **Allergic Reactions:**

Allergic reactions (i.e., pruritus, rash, fever, injection site reaction, bulleous eruption) and skin necrosis have occurred rarely. A few cases of anaphylactoid reactions have been reported.

## **Local Reactions:**

Pain at the injection site, the only non-bleeding event determined to be possibly or probably related to treatment with FRAGMIN and reported at a rate of at least 2% in the group treated with FRAGMIN, was reported in 4.5% of patients treated with FRAGMIN 5000 IU qd vs 11.8% of patients treated with heparin 5000 U bid in the abdominal surgery trials. In the hip replacement trials, pain at injection site was reported in 12% of patients treated with FRAGMIN 5000 IU qd vs 13% of patients treated with heparin 5000 U tid.

## **Ongoing Safety Surveillance:**

Since first international market introduction in 1985, there have been six reports of epidural or spinal hematoma formation with concurrent use of dalteparin sodium and spinal/epidural anesthesia or spinal puncture. Five of the six patients had post-operative indwelling epidural catheters placed for analgesia or received additional drugs affecting hemostasis. The hematomas caused long-term or permanent paralysis (partial or complete) in four of these cases. The sixth patient experienced temporary paraplegia but made a full recovery. Because these events were reported voluntarily from a population of unknown size, estimates of frequency cannot be made.

## **OVERDOSAGE**

### **Symptoms/Treatment:**

An excessive dosage of FRAGMIN Injection may lead to hemorrhagic complications. These may generally be stopped by the slow intravenous injection of protamine sulfate (1% solution), at a dose of 1 mg protamine for every 100 anti-Xa IU of FRAGMIN given. A second infusion of 0.5 mg protamine sulfate per 100 anti-Xa IU of FRAGMIN may be administered if the APTT measured 2 to 4 hours after the first infusion remains prolonged. Even with these additional doses of protamine, the APTT may remain more prolonged than would usually be found following administration of conventional heparin. In all cases, the anti-Factor Xa activity is never completely neutralized (maximum about 60 to 75%).

Particular care should be taken to avoid overdosage with protamine sulfate. Administration of protamine sulfate can cause severe hypotensive and anaphylactoid reactions. Because fatal reactions, often resembling anaphylaxis, have been reported with protamine sulfate, it should be given only when resuscitation techniques and treatment of anaphylactic shock are readily available. For additional information, consult the labeling of Protamine Sulfate Injection, USP, products. A single subcutaneous dose of 100,000 IU/kg of FRAGMIN to mice caused a mortality of 8% (1/12) whereas 50,000 IU/kg was a non-lethal dose. The observed sign was hematoma at the site of injection.

# Fragmin

brand of dalteparin sodium injection

## DOSAGE AND ADMINISTRATION

### Unstable Angina and Non-Q-Wave Myocardial Infarction:

In patients with unstable angina or non-Q-wave myocardial infarction, the recommended dose of FRAGMIN Injection is 120 IU/kg of body weight, but not more than 10,000 IU, subcutaneously (s.c.) every 12 hours with concurrent oral aspirin (75 to 165 mg once daily) therapy. Treatment should be continued until the patient is clinically stabilized. The usual duration of administration is 5 to 8 days. Concurrent aspirin therapy is recommended except when contraindicated.

Table 9 lists the volume of FRAGMIN to be administered for a range of patient weights.

**Table 9**  
**Volume of FRAGMIN to be Administered by Patient Weight**

Patient weight (lb)	<110	110 to 131	132 to 153	154 to 175	176 to 197	≥198
Patient weight (kg)	<50	50 to 59	60 to 69	70 to 79	80 to 89	≥90
Volume of FRAGMIN (mL) <sup>1</sup>	0.55	0.65	0.75	0.90	1.00	1.00

<sup>1</sup> Calculated volume based on the 9.5 mL multiple-dose vial (10,000 anti-Factor Xa IU/mL)

### Hip Replacement Surgery:

Table 10 presents the dosing options for patients undergoing hip replacement surgery. The usual duration of administration is 5 to 10 days after surgery; up to 14 days of treatment with FRAGMIN have been well tolerated in clinical trials.

**Table 10**  
**Dosing Options for Patients Undergoing Hip Replacement Surgery**

Timing of First Dose of FRAGMIN	Dose of FRAGMIN to be Given Subcutaneously			
	10 to 14 Hours Before Surgery	Within 2 Hours Before Surgery	4 to 8 Hours After Surgery <sup>1</sup>	Postoperative Period <sup>2</sup>
Postoperative Start	---	---	2500 IU <sup>3</sup>	5000 IU qd
Preoperative Start - Day of Surgery	---	2500 IU	2500 IU <sup>3</sup>	5000 IU qd
Preoperative Start - Evening Before Surgery <sup>4</sup>	5000 IU	---	5000 IU	5000 IU qd

<sup>1</sup> Or later, if hemostasis has not been achieved.

<sup>2</sup> Up to 14 days of treatment was well tolerated in controlled clinical trials, where the usual duration of treatment was 5 to 10 days postoperatively.

<sup>3</sup> Allow a minimum of 6 hours between this dose and the dose to be given on Postoperative Day 1. Adjust the timing of the dose on Postoperative Day 1 accordingly.

<sup>4</sup> Allow approximately 24 hours between doses.

### Abdominal Surgery:

In patients undergoing abdominal surgery with a risk of thromboembolic complications, the recommended dose of FRAGMIN is 2500 IU administered by s.c. injection once daily, starting 1 to 2 hours prior to surgery and repeated once daily postoperatively. The usual duration of administration is 5 to 10 days.

In patients undergoing abdominal surgery associated with a high risk of thromboembolic complications, such as malignant disorder, the recommended dose of FRAGMIN is 5000 IU s.c. the evening before surgery, then once daily postoperatively. The usual duration of administration is 5 to 10 days. Alternatively, in patients with malignancy, 2500 IU of FRAGMIN can be administered s.c. 1 to 2 hours before surgery followed by 2500 IU s.c. 12 hours later, and then 5000 IU once daily postoperatively. The usual duration of administration is 5 to 10 days.

# Fragmin

## brand of dalteparin sodium injection

Dosage adjustment and routine monitoring of coagulation parameters are not required if the dosage and administration recommendations specified above are followed.

### Administration:

FRAGMIN is administered by subcutaneous injection. It must not be administered by intramuscular injection.

**Subcutaneous injection technique:** Patients should be sitting or lying down and FRAGMIN administered by deep s.c. injection. FRAGMIN may be injected in a U-shape area around the navel, the upper outer side of the thigh or the upper outer quadrangle of the buttock. The injection site should be varied daily. When the area around the navel or the thigh is used, using the thumb and forefinger, you **must** lift up a fold of skin while giving the injection. The entire length of the needle should be inserted at a 45 to 90 degree angle.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

### HOW SUPPLIED

FRAGMIN Injection is available in the following strengths and package sizes:

0.2 mL single-dose prefilled syringe, affixed with a 27-gauge x 1/2 inch needle.

Package of 10:

2500 anti-Factor Xa IU	NDC 0013-2406-91
5000 anti-Factor Xa IU	NDC 0013-2426-91

9.5 mL multiple-dose vial:

10,000 anti-Factor Xa IU/mL	NDC 0013-2436-06
(95,000 anti-Factor Xa IU/vial)	

Store at controlled room temperature 20° to 25°C (68° to 77°F) [see USP].

 only

U.S. Patent 4,303,651

Manufactured for: Pharmacia & Upjohn Company  
Kalamazoo, MI 49001, USA

By: Vetter Pharma-Fertigung  
Ravensburg, Germany  
(prefilled syringes)

Pharmacia & Upjohn N.V./S.A.  
Puurs, Belgium  
(multiple-dose vial)

